

Platyhelminthes / Trematoda

1 order, 7 families, 4 genera, 7 species

Sampling periods: Jan 1978 to June 1979

Werding (1973), without providing data on the sampling, reports on *Maritrema magdalenae*, a new species of trematodes, found in the intestines of the bird *Limnodromus griseus*. Galeano & Romero (1979a, b and 1981a, b) are all based on the same sampling of fish species in the CGSM from Jan 1978 to June 1979. Unfortunately only from Galeano & Romero (1979a) there was a copy available, reporting on six species of Acanthocolpidae, four of them identified to species level. Galeano & Romero (1981a) reported on additional five families of trematodes.

[Table Platyhelminthes / Trematoda](#)

References (in chronological order)

- W73:** Werding, B. (1973). *Maritrema magdalenae* N. SP., ein Trematode von der Isla de Salamanca, Nordkolumbien. Bulletin of Marine and Coastal Research, 7.
- G79a:** Galeano, M. L., & Romero, M. (1979a). Contribución al conocimiento de parásitos de peces de la Ciénaga Grande de Santa Marta: I. Familia Acanthocolpidae (Trematoda: Digenea). An. Inst. Invest. Mar. Punta Betín, 11, 195-217.
- G81a:** Galeano, M. L., & Romero, M. (1981a). Contribución al conocimiento de parásitos de peces en la Ciénaga Grande de Santa Marta, Colombia. II Parte. Boletín Museo del Mar, 40-53.